

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 June 2001 (21.06.2001)

PCT

(10) International Publication Number
WO 01/43787 A2

(51) International Patent Classification⁷: A61L 26/00

(21) International Application Number: PCT/EP00/12661

(22) International Filing Date:
13 December 2000 (13.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
AL99U000010 14 December 1999 (14.12.1999) IT

(71) Applicant and

(72) Inventor: SACCHI, Maria, Cristina [IT/IT]; Via O. Remotti, 4, I-15040 S. Michele (IT).

(72) Inventor; and

(75) Inventor/Applicant (for US only): BELLANDA, Marco [IT/IT]; Via Dante, 81, I-15100 Alessandria (IT).

(74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi S.p.A., Corso di Porta Vittoria, 9, I-20122 Milan (IT).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A PROCESS FOR PREPARING AN AUTOLOGOUS PLATELET GEL AND MEMBRANES THEREOF, AND A KIT FOR CARRYING OUT THIS PROCESS



(57) Abstract: A process is disclosed for preparing an autologous platelet gel and membranes thereof comprising mixing a platelet concentrate with a calcium salt and batroxobin. This process encompassing the use of batroxobin as the gel activator allows to overcome the prior art processes drawbacks connected with the use for the same purpose of human or bovine thrombin. A kit is also described for carrying out this process.

WO 01/43787 A2

A PROCESS FOR PREPARING AN AUTOLOGOUS PLATELET GEL AND MEMBRANES THEREOF, AND A KIT FOR CARRYING OUT THIS PROCESS.

Field of the invention

The present invention relates to a process for preparing an autologous platelet gel
5 and membranes thereof, and a kit for carrying out this process.

Background of the invention

Marx and colleagues have shown that the use of platelet concentrate represents an innovative method to modulate and accelerate wound healing process, as well as bone regenerative processes (Oral surg., Oral Med., Oral Path 1998, 85: 638-
10 646). Therefore these authors elaborated a technique allowing to isolate and precipitate platelets thereby obtaining an autologous platelet concentrate, which is neither toxic nor immunoreactive. This substance is able to enhance the effects of growth factors contained in platelets granules and which are initiators of metabolic pathway involved in wound healing steps.

15 According to Marx's protocol encompassing the use of a cellular separator, the platelet concentrate is prepared by carrying out a venous blood sample of 500 ml, namely an operation which is accomplished by means a venous catheter. The platelet concentrate is subsequently activated with calcium chloride and bovine thrombin to produce a platelet gel which can be combined with either autologous
20 spongy and cortical bone or bone matrix consisting of bio-glasses.

In Italy Marx technique for the activation of a platelet gel cannot be carried out since bovine thrombin is no longer available on the market. In addition this method involving the use of very high amounts of blood requiring specific instrumentation and skilled personnel for preparing platelet concentrate, cannot be realised in a
25 medical surgery so as to allow a routine use of the platelet gel.

In USP5,585,007 and W0 98/11925 methods for preparing platelet gel are disclosed. These methods, although representing an improvement if compared to Marx's protocol, as they necessitate much lower blood amounts (50-60 ml), they require for platelet gel formation the use of bovine or human thrombin.

30 To date human thrombin can only be prepared by DNA recombinant technique. In Italy substances prepared according to this technique can be used only for in vitro experiments, but not for in vivo experiments.

Technical problem

The need was felt of a process for preparing a platelet gel not presenting the prior art processes drawbacks, connected with the use as the activator of human or bovine thrombin.

Summary of the invention

The Applicant has unexpectedly found that it is possible to overcome the aforementioned drawback, by using as the activator batroxobin in place of thrombin.

The present invention therefore relates to a process for preparing a platelet gel comprising mixing a platelet concentrate with an inorganic or organic calcium salt and batroxobin.

A further object of the present invention relates to a kit for carrying out the process according to the present invention which in particular comprises:

- a) a monouse sterile capsule of transparent plastic or glass material, fitted with a screw cap constituted in the upper part by a perforable membrane coated with a suitable sterile coating, said perforable membrane allowing the insertion of the needle of a syringe containing the platelet concentrate, and contemporaneously avoiding the air contact with the inside of the capsule,
- b) two plug injectors surmounting said perforable membrane containing respectively the calcium salt and batroxobin.

Finally further objects of the present invention are:

- the autologous platelet gel obtained with the process according to the present invention.
- a membrane comprising the autologous platelet gel according to the present invention.

Brief description of the drawings

Figure 1 and 2 represent two preferred embodiments of the kit according to the present invention.

In particular in these figures: (3) represents the sterile monouse capsule, (4) the screw cap constituted in the upper part by a perforable membrane (5). (1) and (2) represent the two plug injectors surmounting said perforable membrane (5). Finally (6) represents the syringe containing the platelet concentrate, and (7)

platelet gel formed during the reaction.

Detailed description of the invention

1. Preferred operating conditions of the process according to the present invention

Preferably the inorganic calcium salt used in the process according to the present invention is calcium chloride, whereas when an organic calcium salt is ^{used} ~~used~~ this is preferably calcium gluconate. According to a particularly preferred embodiment, the process is carried out by using calcium gluconate.

The inorganic or organic calcium salt is in the form of an aqueous solution containing it in amounts ranging from $5 \cdot 10^{-5}$ to 0.5 M.

10 According to a preferred embodiment in the process of the present invention, $8 \cdot 10^{-5}$ M (80 μ M) calcium chloride is utilised.

A more preferred embodiment foresees the use of 0,23 M calcium gluconate.

The batroxobin used is preferably in the form of an aqueous solution having 1 thrombin international unit and is sold under the commercial name Botropase by
15 Ravizza Farmaceutici S.p.A.

2. Method for preparing the platelet concentrate

The platelet concentrate used in the process according to the present invention is prepared from whole blood (sample of 40-50ml of venous blood), collected in 20ml syringes, containing 4 ml sodium citrate used as anticoagulant), by a two phases
20 centrifugation, thereby obtaining an intermediate product being platelet rich plasma. This method allows to recover at least 80% of platelets present in whole blood.

In particular this technique encompasses the following steps. Venous blood is collected in 4 tubes and centrifuged at 180 g for 20 minutes. After this treatment
25 two phases are obtained a dark one constituted by precipitated red and white cells at the bottom of the tubes, and a clear one visible in the upper part of the probe, consisting of platelet rich plasma. This plasma is taken and transferred by means of a Pasteur pipette into other 4 tubes and centrifuged at 580 g for 20 minutes. Thanks to this faster centrifugation (580 g versus 180 g) it is possible to obtain
30 platelets sedimentation as a dense small bottom said "pellet". The liquid supernatant consists of platelet poor plasma.

The tubes are emptied, however leaving in each of them a small aliquot (1.5 ml)

of platelet poor plasma necessary to suspend once again the "pellet" thereby obtaining a homogeneous suspension.

Thus proceeding it is possible to obtain a final platelet concentrate (of about 6 ml) deriving from the unification of the platelets suspensions present in the 4 tubes.

- 5 This platelet concentrate after a rest period of about 15 minutes at room temperature, is ready to be used as reactant in the process according to the present invention.

3. Membranes according to the present invention and method for preparing them

- 10 The membrane according to the present invention may contain active principles such as antibacterial agents, disinfectants, and/or sterile excipients etc .

Preferably the membrane according to the present invention essentially consists of the autologous platelet gel according to the present invention.

These membranes are obtained by simply eliminating the liquid in excess.

- 15 For example if the platelet gel preparation occurs in plastic or glass Petri dishes (like those utilised in cellular cultures), it is possible to obtain a thin membrane, as the gel during the activation takes the form of the container.

In this case by eliminating the liquid in excess by means for example of a Pasteur pipette released during the gel formation it is possible to obtain a pliable membrane having a well defined contour.

- 20 The removed liquid in excess consisting of batroxobin and calcium salt is added to the platelet poor plasma coming from the second centrifugation and determines after 20-30 minutes the formation of a further thick platelet gel .

4. The kit according to the present invention

- 25 Figure 1 and 2 represent preferred embodiment of the kit according to the present invention.

- In particular in these figures: (3) represents the sterile monouse capsule, (4) the screw cap constituted in the upper part by a perforable membrane (5). This membrane is coated with a sterile coating normally used in pharmaceutical and diagnostic industry such as aluminum film or another equivalent material. (1) and
30 (2) represent the two plug injectors surmounting said perforable membrane (5) and containing respectively the inorganic or organic calcium salt (more preferably 0.23 M calcium gluconate), and batroxobin. Finally (6) represents the syringe

containing the platelet concentrate, and (7) platelet gel formed during the reaction.

A manual pressure onto the cap injector (1) allows the outlet of calcium gluconate in amounts of from 0.3 to 0.5 ml into the capsule (3). Analogously a manual pressure onto the cap injector (2) allows the outlet of batroxobin in amounts of from 0.3 to 0.5 ml. The capsule (3) reported in Figure 1 has preferably a capacity of from 10 to 20 ml, whereas the same capsule of the kit reported in figure 2 permits to obtain membranes of diameter of from 3 to 9 cm as the capsule may have a diameter ranging from 3.5 to 10.0 cm.

The process according to the present invention carried out with the aforementioned kit encompasses the following steps:

- A) charging a 10 ml syringe (6) with a platelet concentrate amount ranging from 3 to 9 ml;
- B) removing the sterile coating from the perforable membrane (5) and introducing the needle of the syringe into the capsule (3) through the perforable membrane (5);
- C) pressing in rapid succession the cap injectors (1) and (2) containing respectively the calcium gluconate and batroxobin and the piston of the syringe (6) inside the capsule (3);
- D) gently stirring with a rotating movement for about 30 seconds the capsule (3)
- E) unscrewing the screw cap (4), and taking the platelet gel (7) thus formed.

CLAIMS

- 1 1. A process for preparing a platelet gel comprising mixing a platelet concentrate
2 with calcium chloride and batroxobin.
- 1 2. A process for preparing a platelet gel comprising mixing a platelet concentrate
2 with an organic or inorganic calcium salt and batroxobin.
- 1 3. The process according to claim 2 wherein said organic salt is calcium
2 gluconate.
- 1 4. The process according to anyone of claims 1-3 wherein the calcium salt is in
2 the form of an aqueous solution containing it in amounts ranging from $5 \cdot 10^{-5}$ to 0.5
3 M.
- 1 5. The process according to anyone of claims 1, 2 or 4 wherein $8 \cdot 10^{-5}$ M (80 μ M)
2 calcium chloride is utilised.
- 1 6. The process according to anyone of claims 2-4, wherein 0,23 M calcium
2 gluconate is used.
- 3 7. The process according to anyone of claims 1-6, wherein batroxobin has 1
4 thrombin international unit.
- 1 8. The autologous platelet gel obtained with the process according to anyone of
2 claims 1-7.
- 1 9. A membrane comprising the autologous platelet gel according to claim 8,
2 optionally in combination with active principles and/or sterile excipients .
- 1 10. A membrane consisting essentially of the autologous platelet gel according to
2 claim 8.
- 1 11. A process for preparing the membrane according to anyone of claims 9 and
2 10, comprising eliminating the liquid in excess, from the autologous platelet gel
3 according to claim 8.
- 1 12. A kit for carrying out the process according to the present invention which in
2 particular comprises:
 - 3 a) a monouse sterile capsule (3) of transparent plastic or glass material, fitted
4 with a screw cap (4) constituted in the upper part by a perforable membrane
5 (5) coated with a suitable sterile coating removable at the moment of use,
6 said perforable membrane (5) allowing the insertion of the needle of a
7 syringe (6) containing the platelet concentrate, and contemporaneously

- 8 avoiding the air contact with the inside of said capsule (3);
- 9 b) two plug injectors (1) and (2) surmounting said perforable membrane (5)
- 10 containing respectively the calcium salt and batroxobin.
- 1 13. The process according to anyone of claims 1-7 carried out with the kit
- 2 according to claim 12, comprising the following steps:
- 3 A) charging a 10 ml syringe (6) with a platelet concentrate amount ranging
- 4 from 3 to 9 ml;
- 5 B) removing the sterile coating from the perforable membrane (5) and
- 6 introducing the needle of the syringe (6) into the capsule (3) through the
- 7 perforable membrane (5);
- 8 C) pressing in rapid succession the cap injectors (1) and (2) containing
- 9 respectively the calcium gluconate and batroxobin and the piston of the
- 10 syringe (6) inside the capsule (3);
- 11 D) gently stirring with a rotating movement for about 30 seconds the
- 12 capsule (3);
- 13 E) unscrewing the screw cap (4), and taking the platelet gel (7) thus
- 14 formed.

1/2




FIGURE 1

2/2




FIGURE 2